

RESEARCH

Open Access



Integral inequalities under beta function and preinvex type functions

Izhar Ahmad*

*Correspondence:
drizhar@kfupm.edu.sa
Department of Mathematics
and Statistics, King Fahd
University of Petroleum
and Minerals, Dhahran 31261,
Saudi Arabia

Abstract

In the present paper, the notion of P -preinvex function is introduced and new integral inequalities for this kind of function along with beta function are established. The work extends the results appeared in the literature.

Keywords: Euler beta function, Integral inequality, Holder's inequality, P -preinvex function

Mathematics Subject Classification: 33B15, 26B25, 26D15, 26D10

Background

Convexity plays an important role in economics, management science, engineering, finance and optimization theory. Many interesting generalizations and extensions of classical convexity have been used in optimization and mathematical inequalities. Hanson (1981) introduced the concept of invexity. These functions were named invex by Craven (1981) and η -convex by Kaul and Kaur (1980). Weir and Mond (1988) introduced the concept of preinvex function. Later, Mohan and Neogy (1995) presented few properties of preinvex functions. Some refinements of the mathematical inequalities on convex and generalized convex functions have been investigated in Barani et al. (2012), Chalco-Cano et al. (2012), Dragomir (2001), Dragomir and Agarwal (1998), Fok and Vong (2015), Matloka (2014), Muddassar and Bhatti (2013) and Pachpatte (2004).

Let S be a nonempty subset of R^n and let $\eta : S \times S \rightarrow R^n$.

Definition 1 The set $S \subseteq R^n$ is said to be invex with respect to $\eta(u, v)$ if for every $u, v \in S$ and $t \in [0, 1]$,

$$v + t\eta(u, v) \in S.$$

It is obvious that every convex set is invex with respect to $\eta(u, v) = u - v$, but there exist invex sets which are not convex (see Mohan and Neogy 1995).

Definition 2 (Weir and Mond 1988) The function $f : S \rightarrow R$ is said to be preinvex on S with respect to $\eta(u, v)$, if

$$f(v + t\eta(u, v)) \leq (1 - t)f(v) + tf(u)$$

for every $u, v \in S$ and $t \in [0, 1]$,

The Gauss–Jacobi quadrature formula has the following

$$\int_a^b (x - a)^p (b - x)^q f(x) dx = \sum_{k=0}^{\infty} B_{m,k} f(\gamma_k) + R_m^* |f|, \quad (1)$$

for certain $B_{m,k}$, γ_k and rest $R_m^* |f|$ (see Stancu et al. 2002).

Recently, Liu (2014) obtained several integral inequalities for the left hand side of (1) under the following P -convexity:

The function $f : I \rightarrow R$, where $I \subseteq R$ is said to be P -convex on a convex set, if

$$f(tu + (1 - t)v) \leq f(u) + f(v).$$

for every $u, v \in I$ and $t \in [0, 1]$. For the applications of P -convex function and its generalizations, we refer Akdemir and Ozdemir (2010), Barani and Barani (2012), Liu (2013, 2014), Tunc (2013) and Varosanec (2007).

The main purpose of this paper is to introduce the class of P -preinvex function and derive new inequalities for the left hand side of (1) under these assumptions. The presented results generalize the results of Liu (2014) and references cited therein.

New integral inequalities

Definition 3 The function $f : S \rightarrow R$ is said to be P -preinvex on S with respect to $\eta(u, v)$, if

$$f(v + t\eta(u, v)) \leq f(u) + f(v).$$

for every $u, v \in S$ and $t \in [0, 1]$,

Note that every P -convex function (Liu 2014) is a P -preinvex function with respect to $\eta(u, v) = u - v$ for any $t \in [0, 1]$.

Lemma 1 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If f is P -preinvex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\int_a^{a+\eta(b,a)} (x - a)^p (a + \eta(b, a) - x)^q f(x) dx = \eta(b, a)^{p+q+1} \int_0^1 t^p (1 - t)^q f(a + t\eta(b, a)) dt.$$

Proof It is easy to observe that

$$\begin{aligned} \int_a^{a+\eta(b,a)} (x - a)^p (a + \eta(b, a) - x)^q f(x) dx &= \int_0^1 (a + t\eta(b, a) - a)^p (a + \eta(b, a) - a - t\eta(b, a))^q \\ &\quad \times f(a + t\eta(b, a)) dt \\ &= \eta(b, a)^{p+q+1} \int_0^1 t^p (1 - t)^q f(a + t\eta(b, a)) dt \end{aligned}$$

□

The following definition will be used in the sequel.

Definition 4 The beta function is defined for $x, y > 0$ as

$$\beta(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt.$$

Theorem 1 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If $|f|$ is P -preinvex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \leq \eta(b,a)^{p+q+1} \beta(p+1, q+1) (|f(a)| + |f(b)|).$$

Proof Since $|f|$ is P -preinvex function on $[a, a + \eta(b, a)]$, we have

$$|f(a + t\eta(b, a))| \leq |f(a)| + |f(b)|$$

for all $t \in [0, 1]$. By Theorem 1 and P -preinvexity of $|f|$, we get

$$\begin{aligned} \int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx &= \eta(b,a)^{p+q+1} \int_0^1 t^p (1-t)^q |f(a + t\eta(b, a))| dt \\ &\leq \eta(b,a)^{p+q+1} \int_0^1 t^p (1-t)^q (|f(a)| + |f(b)|) dt \\ &= \eta(b,a)^{p+q+1} \beta(p+1, q+1) (|f(a)| + |f(b)|) \\ &\quad \text{(by the definition 4).} \end{aligned}$$

Theorem 2 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If $|f|^{\frac{k}{k-1}}$ is P -preinvex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\begin{aligned} \int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \\ \leq \eta(b,a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left(|f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}} \right)^{\frac{k-1}{k}}. \end{aligned}$$

Proof The P -preinvexity of $|f|^{\frac{k}{k-1}}$ on $[a, a + \eta(b, a)]$ along with Lemma 1, Definition 4 and Hölder inequality imply that

$$\begin{aligned} \int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \\ \leq \eta(b,a)^{p+q+1} \left[\int_0^1 t^{kp} (1-t)^{kq} \right]^{\frac{1}{k}} \left[\int_0^1 |f(a + t\eta(b, a))|^{\frac{k}{k-1}} dt \right]^{\frac{k-1}{k}} \\ \leq \eta(b,a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left[\int_0^1 \left(|f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}} \right) dt \right]^{\frac{k-1}{k}} \\ = \eta(b,a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left(|f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}} \right)^{\frac{k-1}{k}}. \end{aligned}$$

This completes the proof. \square

Theorem 3 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If $|f|^l$ is P -preinvex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \leq \eta(b,a)^{p+q+1} \beta(p+1, q+1) \left(|f(a)|^l + |f(b)|^l \right).$$

Proof The P -preinvexity of $|f|^l$ on $[a, a + \eta(b, a)]$ along with Lemma 1, Definition 4 and Hölder inequality give

$$\begin{aligned} & \int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \\ &= \eta(b,a)^{p+q+1} \int_0^1 [t^p(1-t)^q]^{\frac{l-1}{l}} [t^p(1-t)^q]^{\frac{1}{l}} f(a+t\eta(b,a)) dt \\ &\leq \eta(b,a)^{p+q+1} \left[\int_0^1 t^p(1-t)^q dt \right]^{\frac{l-1}{l}} \left[\int_0^1 t^p(1-t)^q |f(a+t\eta(b,a))|^l dt \right]^{\frac{1}{l}} \\ &\leq \eta(b,a)^{p+q+1} [\beta(p+1, q+1)]^{\frac{l-1}{l}} \left[\left(|f(a)|^l + |f(b)|^l \right) \beta(p+1, q+1) \right]^{\frac{1}{l}} \\ &= \eta(b,a)^{p+q+1} \beta(p+1, q+1) \left(|f(a)|^l + |f(b)|^l \right)^{\frac{1}{l}}. \end{aligned}$$

This completes the proof. \square

Integral inequalities involving prequasi-invex

I state the following theorems as the proof follow on the same lines of the theorems of “[New integral inequalities](#)” section.

Definition 5 (Pinni 1991) The function $f : S \rightarrow R$ is said to be prequasi-invex on S with respect to $\eta(u, v)$, if

$$f(tu + (1-t)v) \leq \max(f(u), f(v))$$

for every $u, v \in S$ and $t \in [0, 1]$.

Theorem 4 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If f is prequasi-invex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$

$$\int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \leq \eta(b,a)^{p+q+1} \beta(p+1, q+1) \max(f(a), f(b)).$$

Theorem 5 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If $|f|$ is prequasi-invex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \leq \eta(b,a)^{p+q+1} \beta(p+1, q+1) \max(|f(a)|, |f(b)|).$$

Theorem 6 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If $|f|^{\frac{k}{k-1}}$ is prequasi-invex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\begin{aligned} & \int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \\ & \leq \eta(b,a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \max\left(|f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}}\right)^{\frac{k-1}{k}}. \end{aligned}$$

Theorem 7 Let $f : S = [a, a + \eta(b, a)] \rightarrow R$ be a continuous function on the interval of real numbers S^0 (the interior of S) with $a < a + \eta(b, a)$. If $|f|^l$ is prequasi-invex function on $[a, a + \eta(b, a)]$, then for some fixed $p, q > 0$,

$$\int_a^{a+\eta(b,a)} (x-a)^p (a+\eta(b,a)-x)^q f(x) dx \leq \eta(b,a)^{p+q+1} \beta(p+1, q+1) \max(|f(a)|^l, |f(b)|^l).$$

Remark 1 If $\eta(b, a) = b - a$ in the theorems of “[Integral inequalities involving pre-quasi-invex](#)” section, then we get the Theorems appeared in Liu (2013).

Conclusion

In this paper, I have introduced the P-preinvex function and used it along with beta function to establish the new integral type inequalities. I also stated the other integral type inequalities under prequasi-invex function. The presented results may be further generalized under weaker convexity assumptions.

Acknowledgements

This research is supported by King Fahd University of Petroleum and Minerals, Saudi Arabia under the Internal Project No. IN131038. The author is thankful to the anonymous referees for their valuable suggestions, which have substantially improved the presentation of the paper.

Competing interests

The author declares that he has no competing interests.

Received: 12 December 2015 Accepted: 15 April 2016

Published online: 26 April 2016

References

- Akdemir AO, Ozdemir ME (2010) Some Hadamard-type inequalities for coordinated P-convex functions and Godunova-Levin functions. AIP Conf Proc 139:7–15
- Barani A, Barani S (2012) Hermite–Hadamard type inequalities for functions when a power of the absolute value of the first derivative is P-convex. Bull Austral Math Soc 86:126–134
- Barani A, Ghazanfari AG, Dragomir SS (2012) Hermite–Hadamard inequality for functions whose derivatives absolute values are preinvex. J Inequal Appl 2012:47
- Craven BD (1981) Invex functions and constrained local minima. Bull Aust Math Soc 24:357–366
- Chalco-Cano Y, Flores-Franulic A, Roman-Flores H (2012) Ostrowski type inequalities for interval-valued functions using generalized Hukuhara derivative. Comput Appl Math 31:457–472
- Dragomir SS (2001) Refinements of the Hermite–Hadamard integral inequality for log-convex function. Aust Math Soc Gaz 28:129–134

- Dragomir SS, Agarwal RP (1998) Two inequalities for differentiable mapping and applications to special mean of real numbers and to trapezoidal formula. *Appl Math Lett* 11:91–95
- Fok H, Vong S (2015) Generalizations of some Hermite–Hadamard-type inequalities. *Indian J Pure Appl Math* 46:359–370
- Kaul RN, Kaur S (1980) Optimality criteria in nonlinear programming involving nonconvex functions. *J Math Anal Appl* 105:104–112
- Hanson MA (1981) On sufficiency of the Kuhn-Tucker conditions. *J Math Anal Appl* 80:545–550
- Liu W (2013) New integral inequalities via (α, m) -convexity and quasi-convexity. *Hacet J Math* 42:289–297
- Liu W (2014) Some Simpson type inequalities for h -convex and (α, m) -convex functions. *J Comput Anal Appl* 16:1005–1012
- Liu W (2014) New integral inequalities involving beta function via P -convexity. *Miskolc Math Notes* 15:585–591
- Matloka M (2014) Inequalities for h -preinvexs. *Appl Math Comput* 234:52–57
- Mohan SR, Neogy SK (1995) On invex sets and preinvex functions. *J Math Anal Appl* 189:901–908
- Muddassar M, Bhatti MI (2013) Some generalizations of Hadamard's-type inequalities through differentiability for s -convex functions. *Indian J Pure Appl Math* 44:131–151
- Pachpatte BG (2004) A note on integral inequalities involving two log-convex functions. *Math Inequal Appl* 7:511–515
- Pinni R (1991) Invexity and generalized convexity. *Optimization* 22:513–525
- Stancu DD, Coman G, Blaga P (2002) *Analiza Numerica si teroia aproximarii*, vol II. Presa Universitara Clujeana, Cluj-Napoca
- Tunc M (2013) On some integral inequalities via h -convexity. *Miskolc Math Notes* 14:1041–1057
- Varosanec S (2007) On h -convexity. *J Math Anal Appl* 326:303–311
- Weir A, Mond B (1988) Preinvex functions in multiple objective optimization. *J Math Anal Appl* 136:29–38

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
